

Name of the medicinal product

Wellness

2. Qualitative and quantitative composition

Each Round Film coated Tablet contains:

- Paracetamol 325.00mg
- Pseudoephedrine hydrochloride 30.00mg
- Dextromethorphan hydrobromide 4.5 mg

Each Oblong Film coated Tablet contains:

- Paracetamol 500.00mg
- Pseudoephedrinehydrochloride 30.00mg
- Diphenhydramine Hydrochloride 25.00 mg

3. Pharmaceutical form

Day Tablet: Orange to Yellowish orange round biconvex unscored Film coated Tablet.

Night Tablet: White to off-white oblong biconvex unscored Film coated Tablet.

4. Clinical particulars

4.1 Therapeutic indications

For the relief of the symptoms associated with colds and influenza.


4.2 Posology and method of administration

Posology

Adults and Children over 12 years

Four tablets should be taken daily:

Dr. Khadija Fathy Alaa



One Round (Orange) tablet to be taken every 4 to 6 hours during the day (no more than three ~~white~~ ^{orange} tablets a day) *Dr. Alaa 3/8/2022*

One Oblong (White) tablet to be taken at night.

Take only one tablet at a time and only at the times of day indicated on the pack. Do not take the nighttime tablets during the day.

Elderly

As for adults (see Pharmacokinetics).

Children

Not recommended for children under 12 years of age.

Method of Administration

For oral use

Hepatic Dysfunction

Caution should be exercised when administering this medicine to patients with severe hepatic impairment.

Renal Dysfunction

Caution should be exercised when administering this medicine to patients with moderate to severe renal impairment.

4.3 Contraindications

Use in individuals with known hypersensitivity to the product or to any of the excipients listed in section 6.1.

This product is contraindicated in individuals with known hypersensitivity to dextromethorphan or to any of the excipients listed in section 6.1.

Dextromethorphan should not be used in patients taking monoamine oxidase inhibitors (MAOIs), or within 14 days of stopping MAOI treatment (see section 4.5). There is a risk of serotonin syndrome with the concomitant use of dextromethorphan and MAOIs and the concomitant use of these medications may cause a rise in blood pressure and hypertensive crisis (see section 4.5)

This product is contraindicated in patients taking serotonin reuptake inhibitors (SSRIs, see section 4.5).

